

BIRKHOFF NORMAL FORMS IN SEMI-CLASSICAL INVERSE PROBLEMS

A. IANTCHENKO, J. SJÖSTRAND, AND M. ZWORSKI

1. INTRODUCTION

The purpose of this note is to apply the recent results on semi-classical trace formulæ [17], and on quantum Birkhoff normal forms for semi-classical Fourier Operators [12] to inverse problems. We show how the *classical Birkhoff normal form* can be recovered from semi-classical spectral invariants. In fact the full quantum Birkhoff normal form of the quantum Hamiltonian near a closed orbit, and infinitesimally with respect to the energy can be recovered. This generalizes recent results of Guillemin [7] and Zelditch [19],[20], [21] obtained in the high energy setting (a special case of semi-classical asymptotics).

We will illustrate the results in a new setting to which they apply. Let $P(h)$ be a semi-classical Schrödinger operator:

$$(1.1) \quad P(h) = -h^2 \Delta + V(x) - E, \quad V \in \mathcal{C}^\infty(\mathbb{R}^{n+1}; \mathbb{R}),$$

with the principal symbol $p(x, \xi) = \xi^2 + V(x) - E$. We make the following assumptions:

$$(1.2) \quad \exists \epsilon > 0, \quad p^{-1}([- \epsilon, \epsilon]) \Subset T^* \mathbb{R}^{n+1}, \quad dp \neq 0 \text{ on } p^{-1}(0).$$

The first assumption guarantees that $P(h)$ has a discrete spectrum near 0, and the second one, that the energy surface $p^{-1}(0)$ is smooth.

The semi-classical inverse problem can be formulated as follows:

What information about the energy surface, $\xi^2 + V(x) = E$, can be recovered from the asymptotics of the spectrum of $-h^2 \Delta + V(x)$ in a small fixed neighbourhood of E , as $h \rightarrow 0$?

This is very natural and can be considered as a mathematical formulation of the general problem arising in spectroscopy. In concrete physical situations, the relation between the Hamiltonian and the spectrum is investigated using *spectroscopic Hamiltonians* which are essentially Hamiltonians in Birkhoff normal forms – see [18] for one of the first uses of Birkhoff normal forms in theoretical chemistry, and [13] for recent applications using experimental spectral data.

The energy surface, $p^{-1}(0) \subset T^* \mathbb{R}^n$ has a natural bicharacteristic foliation, coming from the Hamilton vector field, $H_p = \sum_{j=1}^n \partial_{\xi_j} p \partial_{x_j} - \partial_{x_j} p \partial_{\xi_j}$. Since it describes classical dynamics, its properties are a natural target of a recovery procedure based on quantum information, such as the spectrum. A naïve intuition suggests that the quantum bound states (corresponding to eigenvalues) should correspond to the closed orbits of the classical dynamics. The simplest symplectic invariants associated to such closed orbits, γ , are the actions,

$$S_\gamma = \int_\gamma \xi dx,$$

the eigenvalues of the linear Poincaré map, P_γ , and the *Maslov index*, $\nu_\gamma \in \mathbb{Z}_4$. We recall that P_γ is the differential of the (non-linear) Poincaré map, κ_γ , defined as follows: let U be a hypersurface

in $p^{-1}(0)$, transversal to γ , locally defined near some point on γ . Then U is a symplectic manifold. The symplectic map $\kappa_\gamma : U \rightarrow U$ is defined as the *first return map*:

$$\kappa_\gamma(m) = \exp(T(m)H_p)(m) \in U, \quad \exp(tH_p)(m) \notin U, \quad 0 < t < T(m),$$

where, strictly speaking, the definition may require replacing U by a larger neighbourhood as the range of κ_γ .

The relation between the spectrum and closed orbits is provided by trace formulæ: the Gutzwiller and Balian-Bloch trace formulæ in physics and their mathematical precursor, the Selberg trace formula, and successors, the Duistermaat-Guillemin and the Guillemin-Melrose formulæ (see [17] and references given there). Gutzwiller's formula states that if $\det(I - P_\gamma^k) \neq 0$, for $0 \neq kT_\gamma \in \text{supp } g$, then for $g, \chi \in \mathcal{C}_c^\infty(\mathbb{R})$, $\chi \equiv 1$ near 0, $g \equiv 0$ near 0,

$$(1.3) \quad \text{tr } \hat{g} \left(\frac{P(h)}{h} \right) \chi(P(h)) \sim \sum_{\gamma} \sum_{0 \neq k = -\infty}^{\infty} \frac{e^{ikS_\gamma/h + i\nu_{\gamma,k} \frac{\pi}{2}} T_\gamma}{|\det(P_\gamma^k - I)|^{\frac{1}{2}}} \sum_{j=0}^{\infty} a_{j,\gamma,k}(g) h^j, \quad a_{0,\gamma,k} = g(kT_\gamma).$$

where T_γ is the period of γ , and the γ summation is over all simple closed orbits of H_p on $p^{-1}(0)$. The Maslov index, $\nu_{\gamma,k}$, may in principle depend on the number of times of “going around” the orbit. The coefficients $a_{j,\gamma,k}$ are distributions on $\mathbb{R}$: $\mathcal{C}_c^\infty(\mathbb{R}) \ni g \mapsto a_{j,\gamma,k}$.

It is now standard that under the non-degeneracy and simplicity assumptions, $\det(I - P_\gamma) \neq 0$, $T_\gamma = T_{\gamma'} \Rightarrow \gamma = \gamma'$, we can recover the actions, S_γ , periods, T_γ , Maslov indices, $\nu_{\gamma,k}$, and $|\det(I - P_\gamma^k)|$, for all k . Under stronger non-degeneracy conditions (see (1.4) and (2.9) below), a theorem of Fried [5] (see Proposition 3.2 below) then shows that the eigenvalues of the linear Poincaré map, P_γ , can be recovered.

The higher order coefficients $a_{j,\gamma,k}$ contain further information and they form a family of semi-classical spectral invariants. It turns out that under a stronger version of the non-degeneracy hypothesis, we can recover from them the full infinitesimal information about the non-linear Poincaré map κ_γ . For simplicity we state it here in the case of elliptic orbits only, which is the semi-classical analogue of Guillemin's result [7] – see Sect.5 for the general statement.

Thus suppose that the eigenvalues of $d\kappa_\gamma(w_0)$, $w_0 = U \cap \gamma$, are given by $\exp(\pm i\theta_j)$, $j = 1, \dots, n$, $\theta_j \in (0, \pi)$. We strengthen the non-degeneracy assumption to independence from 2π over rationals:

$$(1.4) \quad \sum_{j=1}^n k_j \theta_j \in 2\pi\mathbb{Z}, \quad k_j \in \mathbb{Z} \implies k_j = 0, \quad j = 1, \dots, n.$$

The now classical normal form theorem of Birkhoff, Lewis and Sternberg says that there exist symplectic coordinates $(x, \xi) \in \mathbb{R}^{2n}$, centered at w_0 , such that in a neighbourhood of w_0 , we have

$$(1.5) \quad \begin{aligned} \kappa_\gamma &= \kappa_0 + \kappa^\flat, \quad \kappa^\flat(x, \xi) = \mathcal{O}((x^2 + \xi^2)^\infty), \\ \kappa_0(\iota, \omega) &= \left(\iota, \omega - \frac{\partial p}{\partial \iota} \right), \quad \iota = (\iota_1, \dots, \iota_n), \quad \omega = (\omega_1, \dots, \omega_n), \\ \iota_j &= \frac{1}{2}(x_j^2 + \xi_j^2), \quad \omega_j = \arg(x_j + i\xi_j), \\ p(\iota) &= p(0) - \sum_{j=1}^n \theta_j \iota_j + \mathcal{O}(\iota^2), \end{aligned}$$

see [7, Theorem 2.1]. The Taylor series of the twist map κ_0 is called the *Birkhoff normal form* of κ . It invariantly describes the infinitesimal properties of the flow near γ . Since we chose θ_j 's in a unique way, $0 < \theta_j < \pi$, the Birkhoff normal form is unique.

Specialized to this case our general result Theorem 4 can be stated as

Theorem 1. *Suppose that $P(h) = -h^2\Delta + V(x) - E$ is a semi-classical Schrödinger operator satisfying (1.2), and that γ is an elliptic closed orbit of H_p on $p^{-1}(0)$, with eigenvalues of P_γ , $e^{\pm i\theta_j}$, satisfying (1.4). Then the leading term and the coefficients $a_{j,\gamma,k}$ in (1.3), determine the Birkhoff normal form of κ_γ .*

As stated above, our motivation for the recovery of the classical Birkhoff normal forms came from an attempt to understand the results of Guillemin [7] and Zelditch [19],[20],[21], in the context of the recent trace formula of Sjöstrand-Zworski [17], and of the (quantum) Birkhoff normal form for semi-classical Fourier integral operator of Iantchenko [11] and Iantchenko-Sjöstrand [12] (see also [16]). Guillemin and Zelditch considered the high energy (or C^∞ singularities) case, corresponding to $h^2\Delta_g - 1$ semi-classically. Guillemin's starting point was his earlier work with Françoise [4], and an observation of Zelditch on the non-commutative residues of (classical) Fourier integral operator [6]. Zelditch's approach also used the non-commutative residue but was more concrete and computational.

In addition to recovering the classical Birkhoff normal form, [7], [19], [20], [21] were concerned with recovering the full quantum Birkhoff normal form of the operator near the closed trajectory. Without giving a definition of the Birkhoff normal form this result follows from

Theorem 2. *Suppose that the assumptions of Theorem 1 are satisfied for potentials V and $\tilde{V}$, with orbits γ and $\tilde{\gamma}$, respectively. If, in the notation of (1.3), $S_\gamma = \tilde{S}_{\tilde{\gamma}}$, $T_\gamma = \tilde{T}_{\tilde{\gamma}}$, and $|\det(I - P_\gamma^k)| = |\det(I - \tilde{P}_{\tilde{\gamma}}^k)|$, $\nu_{\gamma,k} = \tilde{\nu}_{\tilde{\gamma},k}$, for all k , then*

$$\forall j, k, \quad a_{j,\gamma,k} = \tilde{a}_{j,\tilde{\gamma},k} \iff \begin{cases} \text{There exists a unitary } h\text{-Fourier integral operator } F, \\ \text{microlocally defined in a neighbourhood of } \gamma, \\ \text{such that } (-h^2\Delta + \tilde{V})F = F(-h^2\Delta + V), \\ \text{microlocally to infinite order along } \gamma. \end{cases}.$$

The general result follows from Proposition 4.1 and Theorem 4 below, and is given in Corollary 5.2. The definitions of h -Fourier integral operators, and of agreement “to infinite order” are recalled in Sect. 2.

Our approach is a straightforward application of (2.3) and (2.13) quoted below from our earlier papers, and of a result of Fried [5] on the recovery of the eigenvalues of P_γ (see Proposition 3.2 below). The trace formula shows that the traces of powers of the quantum monodromy operator of [17] are spectral invariants. The quantum monodromy operator is a semi-classical operator which quantizes the Poincaré map for a given orbit γ , and once it is put into the Birkhoff normal form using (2.13) the traces of its powers are easy to compute – see Proposition 3.1. The recovery of the classical Birkhoff normal form is then very clear – see Theorem 3.

One of the most striking applications of this approach to inverse problems is the result of Zelditch [21] on the recovery of bi-axial analytic planar domains from the spectrum of the (Dirichlet or Neumann) Laplacian. By applying our general methods we avoid any specific new work involving the boundary value problem – see Theorem 5 and Corollary 5.4. This and other inverse results are presented in Sect.5.

What may seem like an excessive generality in the trace formulæ of [17] (see (2.3) below) and Theorem 4 is motivated by their potential use for the recovery of classical dynamics of *effective Hamiltonians*. One of the most impressive inverse procedures is the application of the *Onsager rules* for determining Fermi surfaces – see for instance [1] for an introductory physics discussion. They can be interpreted in terms of a trace formula involving an effective Hamiltonian coming from the *Peierls substitution* – see [9]. The question which will be investigated elsewhere is: can a more detailed information about the Fermi surface be obtained from higher order terms in the expansions of magnetic susceptibilities of metals.

Throughout the paper we will write

$$u(x, \epsilon) = \sum_{j=0}^{\infty} u_j(x) \epsilon^j,$$

to denote asymptotic expansions as $\epsilon \rightarrow 0$. When no confusion is likely to arise all equalities are meant modulo $\mathcal{O}(\epsilon^\infty)$ where ϵ is the relevant small parameter (mostly h , the mathematical “Planck constant”). We denote by $\mathbb{N}$ the set of natural numbers, $0, 1, 2, \dots$.

ACKNOWLEDGEMENTS. The third author would like to thank the National Science Foundation for partial support under grant DMS-9970614. He would also like to thank Bill Miller and Howard Taylor for references to the chemistry literature.

2. PRELIMINARIES

In this section we will recall the general trace formula of [17] and the classical and quantum Birkhoff normal forms of [12].

We consider X which is either a compact $\mathcal{C}^\infty$ manifold of dimension $n+1$ or $\mathbb{R}^{n+1}$.

We introduce the usual class of semi-classical symbols on X :

$$S^{m,k}(T^*X) = \{a \in \mathcal{C}^\infty(T^*X \times (0, 1]) : |\partial_x^\alpha \partial_\xi^\beta a(x, \xi; h)| \leq C_{\alpha,\beta} h^{-m} \langle \xi \rangle^{k-|\beta|}\},$$

and the corresponding class of pseudodifferential operators, $\Psi_h^{m,k}(X)$, with the quantization and symbol maps:

$$\begin{aligned} \text{Op}_h^w : S^{m,k}(T^*X) &\longrightarrow \Psi_h^{m,k}(X) \\ \sigma_h : \Psi_h^{m,k}(X) &\longrightarrow S^{m,k}(T^*X)/S^{m-1,k-1}(T^*X), \end{aligned}$$

with both maps surjective, and the usual properties

$$\sigma_h(A \circ B) = \sigma_h(A) \sigma_h(B),$$

$$0 \rightarrow \Psi^{m-1,k-1}(X) \hookrightarrow \Psi^{m,k}(X) \xrightarrow{\sigma_h} S^{m,k}(T^*X)/S^{m-1,k-1}(T^*X) \rightarrow 0,$$

a short exact sequence, and

$$\sigma_h \circ \text{Op}_h^w : S^{m,k}(T^*X) \longrightarrow S^{m,k}(T^*X)/S^{m-1,k-1}(T^*X),$$

the natural projection map.

Let $P(z) \in \mathcal{C}^\infty(I_z; \Psi_h^{0,k}(X))$, $I = (-a, a) \subset \mathbb{R}$, be a family of self-adjoint, principal type operators, such that $\Sigma_z = \{m : \sigma(P(z)) = 0\} \subset T^*X$ is compact. We assume that

$$(2.1) \quad \begin{aligned} &\sigma(\partial_z P(z)) \leq -C < 0, \quad \text{near } \Sigma_z, \quad |\sigma(P(z))| \geq C \langle \xi \rangle^k, \\ &\text{for } |\xi| \geq C \text{ when } X \text{ is a compact manifold, and for } |(x, \xi)| \geq C \text{ if } X = \mathbb{R}^{n+1}. \end{aligned}$$

We also assume that for z near 0, the Hamilton vector field, $H_{p(z)}$, $p(z) = \sigma(P(z))$, has a simple closed orbit $\gamma(z) \subset \Sigma_z$ with period $T(z)$, and that $\gamma(z)$ has a neighbourhood Ω such that

$$(2.2) \quad m \in \Omega \text{ and } \exp t H_{p(z)}(m) = m, \quad p(z, m) = 0, \quad 0 < |t| \leq T(z)N + \epsilon, \quad z \in I, \implies m \in \gamma(z),$$

where $T(z)$ is the period of $\gamma(z)$, assumed to depend smoothly on z . We also write $p(z, m)$ for $\sigma(P(z))(m)$. Let $A \in \Psi_h^{0,0}(X)$ be a microlocal cut-off to a sufficiently small neighbourhood of $\gamma(0)$.

Then if $P(z)$ is an almost analytic extension of $P(z)$, $z \in \mathbb{R}$, $\chi \in \mathcal{C}_c^\infty(I)$, $\tilde{\chi} \in \mathcal{C}_c^\infty(\mathbb{C})$, its almost analytic extension, $f \in \mathcal{C}^\infty(\mathbb{R})$, and $\text{supp } \hat{f} \subset (-N(C_p - \epsilon) + C, N(C_p - \epsilon) - C) \setminus \{0\}$, we have,

$$(2.3) \quad \begin{aligned} & \frac{1}{\pi} \text{tr} \int f(z/h) \bar{\partial}_z [\tilde{\chi}(z) \partial_z P(z) P(z)^{-1}] A \mathcal{L}(dz) = \\ & -\frac{1}{2\pi i} \sum_{-1 \neq k = -N-1}^{N-1} \text{tr} \int_{\mathbb{R}} f(z/h) M(z, h)^k \frac{d}{dz} M(z, h) \chi(z) dz + \mathcal{O}(h^\infty), \end{aligned}$$

where $M(z, h)$ is the quantum monodromy operator. The constant $C_p > 0$, in the condition on $\hat{f}$ depends on $p(z)$.

The *quantum monodromy operator*, $M(z, h)$ is defined as follows. For a point on γ , $m_0 \in \gamma$, we can define the microlocal kernel of $P(z)$ at m_0 , to be the set of families $u(h)$, such that $u(h)$ are microlocally defined near m_0 and $P(z)u(h) = \mathcal{O}(h^\infty)$ near m_0 . We denote it by $\ker_{m_0(z)}(P(z))$. Any solution can be continued microlocally along $\gamma(z)$ and we denote the corresponding forward and backward continuation by $I_\pm(z)$ – see [17, §4] for precise definitions. Now, let $m_1 \neq m_0$ be another point on $\gamma(z)$. We then define

$$(2.4) \quad \begin{aligned} I_-(z) \mathcal{M}(z) &= I_+(z), \quad \text{near } m_1, \\ \mathcal{M}(z) : \ker_{m_0(z)}(P(z)) &\longrightarrow \ker_{m_0(z)}(P(z)). \end{aligned}$$

The operator $P(z)$ is assumed to be self-adjoint with respect to some inner product $\langle \bullet, \bullet \rangle$, and we define the *quantum flux* norm on $\ker_{m_0(z)}(P(z))$ as follows: let χ be a microlocal cut-off function supported near γ and equal to one near the part of γ between m_0 and m_1 . We denote by $[P(z), \chi]_+$ the part of the commutator supported in m_0 , and put

$$\langle u, v \rangle_{\text{QF}} \stackrel{\text{def}}{=} \langle [(h/i)P(z), \chi]_+ u, v \rangle, \quad u, v \in \ker_{m_0(z)}(P(z)).$$

It is easy to check that this norm is independent of the choice of χ – see [17, Lemma 4.4]. This independence leads to the unitarity of $\mathcal{M}(z)$:

$$\langle \mathcal{M}(z)u, \mathcal{M}(z)u \rangle_{\text{QF}} = \langle u, u \rangle_{\text{QF}}, \quad u \in \ker_{m_0(z)}(P(z)).$$

For practical reasons we identify $\ker_{m_0(z)}(P(z))$ with $\mathcal{D}'(\mathbb{R}^n)$, microlocally near $(0, 0)$, and choose the identification so that the corresponding monodromy map is unitary (microlocally near $(0, 0)$ where $(0, 0)$ corresponds to the closed orbit intersecting a transversal identified with $T^*\mathbb{R}^n$). This gives

$$M(z, h) : \mathcal{D}'(\mathbb{R}^n) \longrightarrow \mathcal{D}'(\mathbb{R}^n),$$

microlocally defined near $(0, 0)$ and

$M(z, h)$ is a semi-classical Fourier integral operator which quantizes the Poincaré map of $\gamma(z)$. This, and (2.3), (2.6) are the *only* properties of $M(z, h)$ which will be used in this paper.

Basic ideas become clear when one considers the simplest example $P(z) = hD_x - z$, on $\mathbb{S}^1$. In that case $M(z, h) = \exp(2\pi i z/h)$ – see [17, §2] for a careful presentation.

We will now assume that the symbol of $P(z)$ has an asymptotic expansion in powers of h , near Σ_0 :

$$(2.5) \quad (P(z))(x, \xi; h) = \sum_{j=0}^{\infty} h^j (p_j(z))(x, \xi), \quad (x, \xi) \in U, \quad U \text{ a precompact neighbourhood of } \Sigma_0.$$

Under this assumption we have a simple extension of [17, Proposition 7.5]

$$(2.6) \quad \text{tr} M(z, h)^{k-1} h D_z M(z, h) = e^{ikI(z)} \frac{e^{i\nu_k(z)\frac{\pi}{2}} I'(z)}{|d\kappa_{\gamma(z)}^k - 1|^{\frac{1}{2}}} + e^{ikI(z)} \sum_{j=1}^{\infty} a_{j,k} h^j,$$

where $I(z)$ is the classical action, ν_k the Maslov index of the k th iterate of $\gamma(z)$ (it is independent of z for z near 0).

We start by recalling the general statement of the classical *Birkhoff normal form* – see [12, Theorem 1.3, Proposition 4.3] and references there.

Let (W, ω) be a symplectic manifold of dimension $2n$, and let $\kappa : W \rightarrow W$, $\kappa(w_0) = w_0$, be a symplectic transformation, $\kappa^* \omega = \omega$, fixing $w_0 \in W$. In (1.5), the *twist map*, κ_0 , is the Birkhoff normal form of κ ,

$$(2.7) \quad \kappa_0 = \exp H_p,$$

where the Hamiltonian p is in the Birkhoff normal form.

The general case presented in [12] is formulated using (2.7): suppose that the eigenvalues of $d\kappa(w_0)$ are given by

$$(2.8) \quad \begin{aligned} & \lambda_j, \lambda_j^{-1}, \quad j = 1, \dots, n \\ & \lambda_j = \exp \mu_j, \quad \mu_j = \alpha_j + i\beta_j, \quad \alpha_j > 0, \quad 0 < \beta_j \notin 2\pi\mathbb{Z}, \quad 1 \leq j \leq n_{\text{ch}}, \\ & \lambda_{j+n_{\text{ch}}} = \exp \mu_{j+n_{\text{ch}}}, \quad \mu_{j+n_{\text{ch}}} = \overline{\mu_j}, \quad 1 \leq j \leq n_{\text{ch}}, \\ & \lambda_j = \exp \mu_j, \quad \mu_j > 0, \quad 2n_{\text{ch}} + 1 \leq j \leq 2n_{\text{ch}} + n_{\text{rh}}, \\ & \lambda_j = \exp \mu_j, \quad \mu_j \in i(\mathbb{R} \setminus 2\pi\mathbb{Z}), \quad n_{\text{rh}} + 2n_{\text{ch}} + 1 \leq j \leq n. \end{aligned}$$

Here $n_e + n_{\text{rh}} + 2n_{\text{ch}} = n$, and e stands for elliptic, rh for real hyperbolic, and ch, for complex hyperbolic.

We replace the condition in the purely elliptic case (1.4) by the more general condition

$$(2.9) \quad \sum_{j=1}^n k_j \mu_j \in 2\pi i\mathbb{Z}, \quad k_j \in \mathbb{Z} \implies k_j = 0.$$

Then the generalization of (1.5) says that there exist symplectic coordinates $(x, \xi) \in \mathbb{R}^n$ centered at $(0, 0)$

(2.10)

$$\begin{aligned} \kappa &= \exp H_p + \kappa^b, \quad \kappa^b(x, \xi) = \mathcal{O}((x^2 + \xi^2)^\infty), \\ p &= p_0 + r, \quad r(x, \xi) = \mathcal{O}((x^2 + \xi^2)^2), \\ p_0(x, \xi) &= \sum_{j=1}^{n_{\text{ch}}} \alpha_j (x_{2j-1} \xi_{2j-1} + x_{2j} \xi_{2j}) - \beta_j (x_{2j-1} \xi_{2j} - x_{2j} \xi_{2j-1}) \\ &+ \sum_{j=2n_{\text{ch}}+1}^{2n_{\text{ch}}+n_{\text{rh}}} \mu_j x_j \xi_j + \sum_{j=2n_{\text{ch}}+n_{\text{rh}}+1}^n \frac{\mu_j}{2i} (x_j^2 + \xi_j^2) \\ r(x, \xi) &= R(\iota), \quad R(\iota) = \sum_{|\alpha| \geq 2} r_\alpha \iota^\alpha, \\ \iota_j &= \begin{cases} (x_{2j-1} \xi_{2j-1} + x_{2j} \xi_{2j} + i(x_{2j-1} \xi_{2j} - x_{2j} \xi_{2j-1}))/2 & 1 \leq j \leq n_{\text{ch}} \\ \bar{\iota}_{j'} & j' = j - n_{\text{ch}}, \quad n_{\text{ch}} + 1 \leq j \leq 2n_{\text{ch}} \\ x_j \xi_j & 2n_{\text{ch}} + 1 \leq j \leq 2n_{\text{ch}} + n_{\text{rh}} \\ (x_j^2 + \xi_j^2)/(2i) & 2n_{\text{ch}} + n_{\text{rh}} + 1 \leq j \leq n. \end{cases} \end{aligned}$$

Observe that $p_0 = \langle \iota, \mu \rangle$. By the *Birkhoff normal form* of κ we mean p_0 and the asymptotic expansion of r .

We now recall the *quantum Birkhoff normal forms* [11] and [12, Theorem 5.1, (4.11)]. We start with some general facts about h -Fourier integral operators. Let W above be a neighbourhood of $w_0 \in T^*X$, where X is a $\mathcal{C}^\infty$ manifold. To $\kappa : W \rightarrow W$ we can associate a class of operators, microlocally defined near w_0 (see [17, §3] for the precise definition of this notion):

$$I_h^m(X, X, \kappa'),$$

the semi-classical *Fourier integral operators* quantizing κ . One way to define them is by considering oscillatory integrals: we identify w_0 with $(0, 0) \in T^*\mathbb{R}^n$ so that

$$(2.11) \quad \begin{aligned} U \in I_h^m(X, X, \kappa') &\iff Uu(x) = h^{-m-\frac{n+N}{2}} \int \int e^{i\phi(x,y,\theta)/h} a(x,y,\theta;h) u(y) dy d\theta, \\ \phi &\in \mathcal{C}^\infty(\mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}^N), \quad \text{graph}(\kappa) = \pi(C_\phi), \\ C_\phi &= \{(x,y,\theta) : \phi'_\theta(x,y,\theta) = 0\} \ni (x,y,\theta) \xrightarrow{\pi} (x,\phi'_x; y, \phi'_y), \end{aligned}$$

where ϕ is assumed to be a non-degenerate phase function in the sense of Hörmander (see [10, Def. 21.2.5] for a related discussion), that is ϕ is smooth and real, C_ϕ is a smooth manifold of dimension $2n$, and π has an injective differential. The amplitude a is assumed to be supported near $(0, 0, 0)$, and to be a classical symbol in the sense of (2.5).

Before stating the main result of [12] we have to introduce the notion of equivalence of families of operators used in that paper [12, §2, §5]: let $U(z) \in I^0(X, X, \kappa'_z)$ and $\tilde{U}(z) \in I^0(X, X, \tilde{\kappa}'_z)$ be two families of operators. Then

$$(2.12) \quad U \equiv \tilde{U}$$

if $\kappa_0(w_0) = \tilde{\kappa}_0(w_0) = w_0$, the two families of transformations, κ_z and $\tilde{\kappa}_z$, agree to infinite order at w_0 and $z = 0$, and the terms in the asymptotic expansions (in powers of h) of ϕ_z , $\tilde{\phi}_z$, a_z , $\tilde{a}_z$ (with amplitudes as in (2.11)) agree to infinite order at $(0, 0, 0)$ and $z = 0$.

Suppose that $U(z)$ quantizes κ_z and κ_0 satisfies the assumptions of (2.10). Also assume that $U(z)$ is elliptic. Then

$$(2.13) \quad \begin{aligned} U(z) &\equiv V(z)^{-1} e^{iP(z,h)/h} V(z), \\ V(z) \in I^0(X, X, \mathcal{C}'_z) &\text{ is a family of elliptic Fourier integral operators near } ((0, 0), w_0) \\ P(z, h) &= p_0(z) + hp_1(z) + h^2p_2(z) + \cdots, \\ p_j(z) &= p_j(z; I_1, \dots, I_j), \quad I_j = \iota_j^w(x, hD_x), \\ p_0(z) &= \sum_{j=1}^n \mu_j(z) I_j + R(z; I_1, \dots, I_n), \quad R(z; \iota) = \mathcal{O}(\iota^2). \end{aligned}$$

When $U(z)$ is microlocally unitary near w_0 , $P(z, h)$, $V(z)$ can be chosen to be microlocally self-adjoint and unitary, respectively, at $((0, 0), w_0)$. We will call the expansion of $P(z, h)$ at $z = 0$, $\iota = 0$, the *infinitesimal Quantum Birkhoff Normal Form* of $U(z)$ at $z = 0$.

We recall that a Fourier integral operator is elliptic if it is associated to a canonical transformation, that is $\mathcal{C}$ above is the graph of a canonical transformation, $((0, 0), w_0) \in \mathcal{C}$, and that its symbol is elliptic, that is the leading term in the asymptotic expansion of the amplitude in (2.11) does not vanish.

3. RECOVERING BIRKHOFF NORMAL FORMS FROM TRACES

From [17, Lemma 3.2, Proposition 7.3] we know that if $U(z) \in I_h^0(X, X, \kappa(z)')$, is a family of h -Fourier integral operators, and that the canonical relations $\kappa(z)^k$, $k \neq 0$, have only one non-degenerate fixed point then

$$(3.1) \quad \text{tr } U(z)^k = e^{ikS(z)/h} (a_{0k}(z) + a_{1k}(z)h + \cdots + a_{lk}(z)h^l + \mathcal{O}(h^{l+1})) .$$

In this section we will prove the following

Theorem 3. *Let $U(z) \in I_h^0(X, X, \kappa(z)')$ be a family of h -Fourier integral operator associated to smooth locally defined canonical transformations, $\kappa(z) : T^*X \rightarrow T^*X$. Suppose that κ_0 has a unique fixed point, w_0 , and that the eigenvalues of $d\kappa_0(w_0)$ given in (2.8) satisfy (2.9). Then the Taylor series of the coefficients $a_{lk}(z)$ in (3.1) determine the infinitesimal quantum Birkhoff normal form of $U(z)$, $P(z, h)$ in (2.13). In particular, the Birkhoff normal form of κ_0 at $z = 0$, $p(z)$, in (2.10), is determined.*

Our proof of Theorem 3 is based on

Proposition 3.1. *Suppose that*

$$(3.2) \quad G(\iota; h) = \langle \iota, \mu \rangle + F(\iota, h), \quad F(\iota, h) = F_0(\iota) + hF_1(\iota) + \cdots, \quad e^{\mu_j} \neq 1, \quad F_0(\iota) = \mathcal{O}(\iota^2),$$

where we consider $F(\iota, h)$ as a formal power series in h , and each $F_j(\iota)$, a formal power series in ι , and where we suppressed the dependence on μ in higher order terms.

If $I_j(h) = \iota_j^w(x, hD_x)$, then

$$(3.3) \quad \text{tr } e^{-iG(I(h); h)/h} = e^{-iF(ih\partial_\mu; h)/h} \left(\prod_{j=1}^n \frac{1}{2 \sinh(\mu_j/2)} \right) .$$

Proof. When $G(\iota; h) = \langle \iota, \mu \rangle$ this follows from computations in each of the cases given in (2.10):

$$\text{tr } e^{-i\langle I(h), \mu \rangle/h} = \text{tr } e^{-i\langle I(1), \mu \rangle},$$

where we can separate the trace into separate blocks:

- *elliptic blocks:*

$$\text{tr } e^{-\mu(D_x^2 + x^2)/2} = \sum_{k=0}^{\infty} e^{-\mu(n + \frac{1}{2})} = \frac{1}{2 \sinh \frac{\mu}{2}},$$

where the formal computation ($\mu \in i\mathbb{R}_+$) can be justified by considering the limit $\epsilon \rightarrow 0$, with μ replaced by $\mu + \epsilon$.

- *real hyperbolic blocks:*

$$\text{tr } e^{-i\mu(xD_x + D_x x)/2} = \text{tr } e^{-\mu(x\partial_x + \frac{1}{2})} = \text{tr} \left(u \mapsto e^{-\mu/2} u(e^{-\mu} \bullet) \right),$$

Formally,

$$\text{tr } e^{-\mu(x\partial_x + \frac{1}{2})} = e^{-\mu/2} \int \delta_0((e^{-\mu} - 1)x) dx = \frac{1}{2 \sinh \frac{\mu}{2}},$$

and this can be justified by approximating δ_0 by a sequence of smooth compactly supported functions.

- *complex hyperbolic blocks*: here we have to work in two dimensions, $\mu_1 = \alpha + i\beta$, $\mu_2 = \bar{\mu}_1 = \alpha - i\beta$, and the corresponding ι_1 and ι_2 (see the last two lines in (2.10)) are $\iota_1 = z \cdot \zeta$, $\iota_2 = w \cdot \omega$, where $z = (x_1 - ix_2)/\sqrt{2}$, $w = \bar{z}$ and $\zeta = (\xi_1 + i\xi_2)/\sqrt{2}$, $\omega = (\xi_1 - i\xi_2)/\sqrt{2}$, the corresponding dual coordinates, $\omega = \bar{\zeta}$. With $I_j = \iota_j^w(x, D_x)$, $j = 1, 2$, we have

$$\text{tr } e^{-i\mu_1 I_1 - i\mu_2 I_2} = \text{tr } e^{-\mu_1(z\partial_z + \frac{1}{2}) - \bar{\mu}_1(\bar{z}\partial_{\bar{z}} + \frac{1}{2})} = \text{tr } \left(u(\bullet, \bullet) \mapsto e^{-\mu_1/2 - \bar{\mu}_1/2} u(e^{-\mu_1} \bullet, e^{-\bar{\mu}_1} \bullet) \right).$$

The last operator can be written as a contour integral,

$$e^{-\mu_1/2 - \bar{\mu}_1/2} u(e^{-\mu_1}, e^{-\bar{\mu}_1} \bar{z}) = \frac{1}{(2\pi)^2} \iiint e^{i((e^{-\mu_1} z - \bar{z})\tilde{\zeta} + (e^{-\bar{\mu}_1} \bar{z} - \tilde{w})\tilde{\omega})} u(\tilde{z}, \tilde{w}) d\tilde{z} d\tilde{w} d\tilde{\zeta} d\tilde{\omega},$$

with the integration contour given by $\tilde{w} = \bar{\tilde{z}}$, $\tilde{\omega} = \bar{\tilde{\zeta}}$.

Formal stationary phase argument applied when taking the trace gives

$$\prod_{j=1}^2 \frac{1}{2 \sinh \frac{\mu_j}{2}}.$$

Once we understand the contribution of the leading term the following formal argument essentially gives (3.3). Since $I(h)$ is unitarily equivalent to $hI(1)$ we see that

$$\text{tr } e^{-iG(I(h);h)/h} = \text{tr } e^{-iG(hI(1);h)/h}.$$

On the other hand, $I(1) = \partial_\mu \langle I(1), \mu \rangle$, and hence

$$e^{-iG(hI(1);h)/h} = e^{-iF(hI(1);h)/h} e^{-i\langle I(1), \mu \rangle} = e^{-iF(ih\partial_\mu;h)/h} e^{-i\langle I(1), \mu \rangle}.$$

By moving differentiation outside of the trace we obtain the expansion (3.3). $\square$

We also need a result of Fried [5] used in the same context in [7]:

Proposition 3.2. *If the assumption (2.9) holds then the set*

$$a_0^k \prod_{j=1}^n \sinh(k\mu_j/2), \quad k \in \mathbb{N},$$

determines $\exp \mu_j$'s and a_0 uniquely. $\square$

Proposition 3.2 and an argument based on Kronecker's Theorem give the following inverse result crucial in the proof of Theorem 3 (see [7, §8], [20, §6]) for different original approaches:

Lemma 3.3. *Suppose that μ_j satisfy (2.9). Then for any polynomial, $p(\xi) = \sum_{|\alpha| \leq M} a_\alpha \xi^\alpha$, the coefficients a_α can be recovered from the asymptotics of*

$$(3.4) \quad p(ik^{-1}\partial_\mu) \prod_{j=1}^n \frac{1}{2 \sinh(k\mu_j/2)}, \quad k \rightarrow \infty.$$

Proof. Using the argument of the proof of Proposition 3.1 we can rewrite (3.4) with $e_0 = (1, \dots, 1)$:

$$(3.5) \quad \begin{aligned} p(ik^{-1}\partial_\mu) \prod_{j=1}^n \frac{1}{2 \sinh(k\mu_j/2)} &= p(ik^{-1}\partial_\mu) \sum_{m \in \mathbb{N}^n} e^{-\langle m + e_0/2, k\mu \rangle} \\ &= \sum_{m \in \mathbb{N}^n} p((m + e_0/2)/i) e^{-\langle m + e_0/2, k\mu \rangle}. \end{aligned}$$

To justify this expression we will consider it as a distribution obtained as a boundary value of a holomorphic function. We introduce new complex variables $z \in \Omega_1 = D(0, 1)^{n_e}$ and $w \in \Omega_2 =$

$(D(0, 1) \setminus 0)^{n_{\text{ch}}}$ in a polydisc and a punctured polydisc respectively: at the boundary, $z_j = e^{i\mu_j/2}$, and $w_j = e^{i\beta_j + n_e + n_{\text{rh}}/2}$. We then write

$$\begin{aligned} m &= (m'_{\text{ch}}, m''_{\text{ch}}, m_{\text{rh}}, m_e) \in \mathbb{N}^n, \quad e_0 = (e_0^{\text{ch}}, e_0^{\text{ch}}, e_0^{\text{rh}}, e_0^e), \\ \mu &= (\mu_{\text{ch}}, \bar{\mu}_{\text{ch}}, \mu_{\text{rh}}, \mu_e), \quad \mu_{\text{ch}} = \alpha + i\beta, \quad \mu_{\text{rh}} \in \mathbb{R}_+^{n_{\text{rh}}}, \quad \mu_e \in (i\mathbb{R})^{n_e}, \\ \langle m + e_0/2, \mu \rangle &= \langle n_2 + e_0^{\text{ch}}, \alpha \rangle + \langle n_1 + e_0^{\text{rh}}/2, \mu_{\text{rh}} \rangle + i\langle l_1 + e^e/2, \mu_e/i \rangle + i\langle l_2, \beta \rangle, \\ n_2 &= m'_{\text{ch}} + m''_{\text{ch}}, \quad l_2 = m'_{\text{ch}} - m''_{\text{ch}}, \quad n_1 = m_{\text{rh}}, \quad l_1 = m_e. \end{aligned}$$

Here $m \in \mathbb{N}^n$ is arbitrary, and

$$n_1 \in \mathbb{N}^{n_{\text{rh}}}, \quad l_1 \in \mathbb{N}^{n_e}, \quad n_2 \in \mathbb{N}^{n_{\text{ch}}}, \quad l_2 \in \mathbb{Z}^{n_{\text{ch}}},$$

with the constraint that

$$\frac{n_2 \pm l_2}{2} \in \mathbb{N}^{n_{\text{ch}}},$$

which implies that $|l_2| \leq n_2$, component wise. We then put

$$q(n, l) = p((m + e_0/2)/i), \quad z = e^{-i\mu_e} \in \mathbb{C}^{n_e}, \quad w = e^{-i\beta/2} \in \mathbb{C}^{n_{\text{ch}}},$$

(component-wise in the natural sense). In this notation (3.5) becomes

$$(3.6) \quad \sum_{n \in \mathbb{N}^{n_{\text{rh}}} + n_{\text{ch}}} e^{-k(\langle n_1 + e_0^{\text{rh}}/2, \mu_{\text{rh}} \rangle + \langle n_2 + e_0^{\text{ch}}/2, \alpha \rangle)} F_n(z^k, w^k),$$

where

$$F_n(z, w) = \sum_{\substack{l_1 \in \mathbb{N}^{n_e}, \quad l_2 \in \mathbb{Z}^{n_{\text{ch}}} \\ \frac{n_2 \pm l_2}{2} \in \mathbb{N}^{n_{\text{ch}}}}} q(n, l) z^{2l_1 + e_0^e} w^{2l_2},$$

$(z, w) \in \Omega_1 \times \Omega_2 \subset \mathbb{C}^{n_e} \times \mathbb{C}^{n_{\text{ch}}}$. Here we note that for any fixed n , $q(n, l)$ is non-zero for finitely many l_2 's and hence $F_n(z, w)$ is holomorphic in the punctured polydisc. Our task is to recover the coefficients $q(n, l)$ from asymptotics as $k \rightarrow 0$. That amounts to recovering the holomorphic functions F_n . We observe that (3.5) shows that the distributional boundary values F_n 's are smooth away from $z_j^4 = 1$.

An application of Kronecker's theorem¹ shows that for any

$$(x, y) \in (\partial D(0, 1))^{n_e} \times (\partial D(0, 1))^{n_{\text{ch}}},$$

there exist a sequence $k_r \rightarrow \infty$ such that $(z^{k_r}, w^{k_r}) \rightarrow (x, y)$.

From (2.9) we deduce that

$$\sum_{i=1}^{n_{\text{rh}} + n_{\text{ch}}} k_j \operatorname{Re} \mu_j = 0 \implies k_j = 0.$$

Hence by choosing sequences of $k \rightarrow \infty$, we can determine the coefficients, $F_n(x, y)$ in the first sum (3.6) (the terms have different rates of exponential decay). From that we determine the holomorphic functions F_n , and the coefficients, $q(n, l)$. $\square$

Proof of Theorem 3. Applying [12, Theorem 3.2] and [12, Theorem 4.4] we see that, up to $\mathcal{O}(h^\infty)$ errors, we can write

$$\operatorname{tr} U(z)^k = \operatorname{tr} e^{-kiG(I(h), z; h)/h},$$

¹which says that if $\alpha \in \mathbb{Z}^n$ satisfies $\langle \alpha, k \rangle \in 2\pi\mathbb{Z}$, $k \in \mathbb{Z}^n$, $\implies k = 0$, then for any $y \in \mathbb{R}^n$ and $\epsilon > 0$, there exists $N \in \mathbb{N}$ and $K \in \mathbb{Z}^n$, such that $|y - N\alpha - 2\pi K| < \epsilon$. Its use is justified by (2.9).

where G is as in Proposition 3.1. Hence,

$$\mathrm{tr} U(z)^k = e^{-ikF(ikh^{-1}\partial_\mu, z; h)/h} \mathrm{tr} e^{-i\langle I(1), k\mu \rangle} \Big|_{\mu=\mu(z)}.$$

Recovering the Birkhoff normal form of κ means recovering the Taylor series of $\langle \iota, \mu(z) \rangle + F_0(\iota, z)$ from the trace. We will actually recover the full infinitesimal *quantum* Birkhoff normal form, $\langle \iota, \mu(z) \rangle + F(\iota, z; h)$.

We write

$$\begin{aligned} \frac{F(hy, z; h)}{h} &= \sum_{j=0}^{\infty} h^j f_j(z, y), \quad f_j(z, y) = \sum_{l=0}^{j+1} \sum_{|\alpha|=j+1-l} \frac{1}{\alpha!} \partial^\alpha F_l(z, 0) y^\alpha, \\ f_j(z, y) &= \sum_{k=0}^{\infty} f_{jk}(y) z^k. \end{aligned}$$

The polynomials $f_{jk}(y)$ clearly determine the formal series $F(\iota, z; h)$.

Since

$$\begin{aligned} \exp(-ikF(hik^{-1}\partial_\mu, z; h)/h) &= \exp(-ik \sum_{j=0}^{\infty} h^j f_j(z, ik^{-1}\partial_\mu)) = \\ (3.7) \quad e^{-ikf_0(z)} &\left(1 - i \sum_{j=1}^{\infty} h^j k f_j(z, ik^{-1}\partial_\mu) - \frac{1}{2!} \left(\sum_{j=1}^{\infty} h^j k f_j(z, ik^{-1}\partial_\mu) \right)^2 + \dots \right), \end{aligned}$$

we see from (3.3), that the coefficient of h^0 in the expansion of the trace is given by

$$\begin{aligned} e^{-ikf_0} &\left(\left(1 - i \sum_{m=1}^{\infty} z^m k f_{0m} - \frac{1}{2!} \left(\sum_{m=1}^{\infty} z^m k f_{0m} \right)^2 + \dots \right) \times \right. \\ &\left. \left(1 - i \sum_{j=1}^{\infty} z^j k \langle ik^{-1}\partial_\mu, \mu^{(j)}(0) \rangle - \frac{1}{2!} \left(\sum_{j=1}^{\infty} z^j k \langle ik^{-1}\partial_\mu, \mu^{(j)}(0) \rangle \right)^2 + \dots \right) \right) \mathrm{tr} e^{-i\langle I(1), k\mu \rangle} \Big|_{\mu=\mu(0)}, \end{aligned}$$

where we expanded $\mu(z)$ into its Taylor series and then used the same argument as in the proof of Proposition 3.1. We apply Proposition 3.2 to the constant term to recover $\exp if_{00}$ and $\mu(0)$. For $j > 0$, the coefficient of z^j in the expression above is given by

$$-ike^{-ikf_0} (f_{0j} + \langle k^{-1}i\partial_\mu, \mu^{(j)}(0) \rangle + H_{j,k}(ik^{-1}\partial_\mu)) \left(\prod_{j=1}^n \frac{1}{2 \sinh(k\mu_j/2)} \right) \Big|_{\mu=\mu(0)},$$

where $H_{j,k}$ is constructed from f_{0l} 's and $\mu^{(p)}(0)$ with $l, p < m$. Hence Lemma 3.3 shows that f_{0j} 's and $\mu^{(j)}(0)$ can be successively recovered.

We now move to higher powers of h , with the procedure being essentially the same: we see from (3.7) that the coefficient of h^m in the expansion of the trace is given by

$$e^{-ikf_0(z)} (-ikf_m(ik^{-1}\partial_\mu, z) + G_{m,k}(ik^{-1}\partial_\mu, z)) \left(\prod_{j=1}^n \frac{1}{2 \sinh(k\mu_j/2)} \right) \Big|_{\mu=\mu(z)},$$

where $G_{m,k}$ is constructed from f_l 's with $l < m$, and we already know the z -expansions of $f_0(z)$ and $\mu(z)$. Replacing f_m by its expansion in z , reduces the problem to recovering the polynomials

$f_{ml}(y)$ from

$$f_{ml}(ik^{-1}\partial_\mu) \prod_{j=1}^n \frac{1}{2 \sinh(k\mu_j/2)}, \quad k \in \mathbb{N},$$

and that follows again from Lemma 3.3. This shows that we recovered the full expansion (in ι , z , and h) of $G(\iota, z; h)$, that is the infinitesimal quantum Birkhoff normal form at $z = 0$. $\square$

4. THE MONODROMY OPERATOR DETERMINES P NEAR γ .

The trace formula (2.6) and the results of Sect.3 will be used to recover the infinitesimal quantum Birkhoff normal form of the quantum monodromy operator $M(z)$ at $z = 0$. In this section we will discuss the recovery of the full Hamiltonian $P(z)$, infinitesimally at $z = 0$, from $M(z)$. Since our information comes from the left hand side of the trace formula (2.6), this recovery can only be possible up to conjugation by families of elliptic h -Fourier integral operators, and up to multiplication by families of elliptic h -pseudo-differential operators. The assumptions (2.1), the implicit function theorem, and the usual symbolic iteration, imply that

$$P(z) = A(z)^*(P - z)A(z),$$

with $A(z) \in \Psi_h^{0,k/2}(X)$ elliptic near $\gamma(0)$, and $P \in \Psi_h^{0,0}(X)$ is self-adjoint. Replacing $P(z)$ by $P - z$ in (2.6) changes the trace by $\mathcal{O}(h^\infty)$. Hence we would like to recover P infinitesimally near γ , on the energy surface $p^{-1}(0)$, $p = \sigma(P)$.

Thus in this section, let P be a self-adjoint h -pseudodifferential operator with leading symbol p . Let γ be a simple closed H_p -trajectory of period $T_0 > 0$. Recall from [17, §4] that the quantum time, Q , is an h -pseudodifferential operator microlocally defined in a suitable neighbourhood $W_+ = \text{neigh}(m_0)$ of some fixed point $m_0 \in \gamma$, such that

$$\frac{i}{h}[P, Q] = Id.$$

After conjugation by a unitary Fourier integral operator, we may assume that

$$(4.1) \quad P = hD_{x_n}, \quad Q = x_n.$$

Let $K(z) : \mathcal{D}'(\mathbf{R}^n) \rightarrow \mathcal{D}'(X)$ be a microlocally defined Fourier integral operator with

$$(4.2) \quad (P - z)K(z) = 0, \quad \frac{i}{h}K(z)^*[P, \chi]_{W_+}K(z) = Id,$$

where $\chi \in C^\infty(W_+)$ is equal to 0 near the incoming part of γ and equal to 1 near the outgoing part, and we let χ also denote a corresponding h -quantization. In the case of (4.1), we can take

$$(4.3) \quad K(z)u(x) = e^{izx_n/h}u(x').$$

It follows from the second part of (4.2), that

$$K(z)^*f'(Q)K(z) = Id,$$

for $f \in C^\infty(\mathbf{R})$, $= 0$ for large negative t and $= 1$ for large positive t . Indeed, $e^{itP/h}Qe^{-itP/h} = Q + t$ and

$$\begin{aligned} \frac{i}{h}[P, f(Q)] &= (\partial_t)_{t=0}(e^{itP/h}f(Q)e^{-itP/h}) = (\partial_t)_{t=0}f(e^{itP/h}Qe^{-itP/h}) \\ &= (\partial_t)_{t=0}f(Q + t) = f'(Q). \end{aligned}$$

Notice that $f' \in C_0^\infty(\mathbf{R})$, $\int f'(q) dq = 1$. More generally,

$$(4.4) \quad K(z)^* g(Q) K(z) = \int g(q) dq \cdot Id$$

for $g \in C_0^\infty(\mathbf{R})$. As in [17, Proposition 4.3] we can choose $K(z)$ so that

$$(4.5) \quad K(z) = e^{i(z-w)Q/h} K(w),$$

since this is fulfilled when (4.1), (4.3) hold (note that for $P(z) = P - z$, $U(z, w) = \exp(i(z-w)Q)$, in [17, (4.4)]).

Consider

$$\pi(z) \stackrel{\text{def}}{=} \frac{1}{2\pi h} K(z) K(z)^*.$$

Notice that this definition is independent of the choice of $K(z)$ satisfying (4.2).

We have the properties:

$$(4.6) \quad \pi(z) = \pi(z)^*,$$

$$(4.7) \quad P\pi(z) = \pi(z)P = z\pi(z),$$

$$(4.8) \quad \int \pi(z) dz = Id,$$

where for the understanding of (4.8), we remark that if $\phi \in C_0^\infty(\mathbf{R})$ is independent of h , then $\int \phi(z)\pi(z) dz$ is a well defined h -pseudodifferential operator of order 0, and that we can always choose ϕ to be equal to 1 in some interval corresponding to the region of phase space where we wish to establish (4.8).

The relations (4.6), (4.7) are obvious, and so is (4.8) if we work in the representation (4.1), (4.3).

We also give a direct proof of (4.8) which does not involve conjugation to the model: Let $g \in C_0^\infty(\mathbf{R})$ and consider

$$K(z)^* g(Q) K(w),$$

where we will assume (4.5). Then we get

$$(4.9) \quad K(z)^* g(Q) K(w) = K^*(z) g(Q) e^{-i(z-w)Q/h} K(z) = (\mathcal{F}_h g)(z-w) \cdot Id,$$

where $\mathcal{F}_h g(z) = \int e^{-iqz/h} g(q) dq$ is the h -Fourier transform. Here the second equality follows from (4.4).

Consider the microlocally defined h -pseudodifferential operator:

$$A = \int \pi(z) dz.$$

A is an elliptic pseudodifferential operator of order 0, and we claim that

$$(4.10) \quad A^2 = A.$$

Since the statement is microlocal, it suffices to show that

$$(4.11) \quad Ag(Q)A = Ag(Q),$$

$g \in C_0^\infty$ (and then choose $g(Q) = 1$ in the region of interest). To get (4.11), we first compute:

$$\begin{aligned} \pi(z)g(Q)\pi(w) &= \frac{1}{(2\pi h)^2} K(z)K(z)^*g(Q)K(w)K(w)^* \\ &= \frac{1}{(2\pi h)^2} K(z)K(z)^*(\mathcal{F}_h g)(z-w)e^{i(z-w)Q/h}, \end{aligned}$$

where we used (4.9), (4.4). Integrating with respect to w , we get

$$\pi(z)g(Q)A = \pi(z)g(Q),$$

which gives (4.10) after integration in z .

Rewriting (4.10) as $A(A - 1) = 0$ and using that A is elliptic, we get (4.8).

Remark. Using (4.3) (or a direct argument), one can check that

$$\pi(z) = \frac{1}{2\pi h} \int e^{it(P-z)/h} dt,$$

where the integral to the right is microlocally well-defined by restricting the integration to a suitable finite interval.

Let $P, \tilde{P}$ be two h -pseudodifferential operators with simple closed trajectories

$$\gamma \subset p^{-1}(0), \quad \tilde{\gamma} \subset \tilde{p}^{-1}(0)$$

with the same period T_0 . Let $m_0 \in \gamma$, $\tilde{m}_0 \in \tilde{\gamma}$ and assume that the corresponding monodromy operators

$$\begin{aligned} \mathcal{M}(z) &: \ker_{m_0}(P - z) \longrightarrow \ker_{m_0}(P - z), \\ \tilde{\mathcal{M}}(z) &: \ker_{\tilde{m}_0}(\tilde{P} - z) \longrightarrow \ker_{\tilde{m}_0}(\tilde{P} - z), \end{aligned}$$

are conjugated:

$$(4.12) \quad \mathcal{U}(z)^{-1} \tilde{\mathcal{M}}(z) \mathcal{U}(z) = \mathcal{M}(z),$$

where $\mathcal{U}(z) : \ker_{\tilde{m}_0}(\tilde{P} - z) \longrightarrow \ker_{m_0}(P - z)$ is a unitary Fourier integral operator depending smoothly on z .

Identifying $P, \tilde{P}$ with hD_{x_n} , we see that

$$F \stackrel{\text{def}}{=} \int \mathcal{U}(z) \pi(z) dz$$

is an elliptic unitary Fourier integral operator with

$$F|_{\ker_{m_0}(P-z)} = \mathcal{U}(z), \quad \tilde{P}F = FP.$$

We can identify $P, \tilde{P}$ with hD_{x_n} in such a way that F becomes the identity operator. With this identification, we take $K(z) = \tilde{K}(z)$ of the form (4.3). Without using the identification, this means that $\tilde{K}(z) = FK(z)$ and

$$(4.13) \quad F = \frac{1}{2\pi h} \int \tilde{K}(z) K(z)^* dz.$$

Also,

$$(4.14) \quad \tilde{M}(z) = M(z).$$

Let $K_f(z), \tilde{K}_f(z); K_b(z), \tilde{K}_b(z)$ be the forward and backward extensions of $K; \tilde{K}$ as in [17, §5].

If u is microlocally concentrated near $m = \exp(tH_p)(m_0)$, $-\epsilon \leq t \leq T_0 - 2\epsilon$, where T_0 is the common period of $\gamma, \tilde{\gamma}$, we define $F_f u$, microlocally concentrated to a neighbourhood of $\exp(tH_{\tilde{p}}(\tilde{m}_0))$, by

$$(4.15) \quad F_f u = e^{-it\tilde{P}/h} F e^{itP/h} u.$$

Since F intertwines $\tilde{P}$ and P , this definition is invariant under small variations of t and we see that F_f is a unitary Fourier integral operator with canonical transformation, given by

$$\kappa_{F_f}(m) = \exp(tH_{\tilde{p}}) \circ \kappa_F \circ \exp(-tH_p)(m),$$

for m in a neighbourhood of $\exp(tH_p)(m_0)$, and where $\kappa_F : \text{neigh}(m_0) \rightarrow \text{neigh}(\tilde{m}_0)$ is the canonical transformation associated to F .

We have

$$F_f u = \frac{1}{2\pi h} \int \tilde{K}_f(z) K_f(z)^* dz.$$

In fact, by (4.13), (4.15):

$$\begin{aligned} F_f &= \frac{1}{2\pi h} \int e^{-it\tilde{P}/h} \tilde{K}_f(z) K_f(z)^* e^{itP/h} dz \\ &= \frac{1}{2\pi h} \int e^{-itz/h} \tilde{K}_f(z) K_f(z)^* e^{itz/h} dz = \frac{1}{2\pi h} \int \tilde{K}_f(z) K_f(z)^* dz. \end{aligned}$$

Similarly, if u is microlocally concentrated near $m = \exp(tH_p)(m_0)$, $-T_0 + 2\epsilon \leq t \leq \epsilon$, we define $F_b u$ to be the right hand side of (4.15) (now with t in a different interval) and get

$$F_b u = \frac{1}{2\pi h} \int \tilde{K}_b(z) K_b(z)^* dz.$$

If m belongs to the overlap region, where both the forward and the backward extensions of F are defined, then for u concentrated to a neighbourhood of m , we have $F_f u = F_b u$:

$$F_f u = \frac{1}{2\pi h} \int \tilde{K}_f(z) K_f(z)^* u dz = \frac{1}{2\pi h} \int \tilde{K}_b(z) \tilde{M}(z) M(z)^* K_b(z)^* u dz = F_b u,$$

since $\tilde{M}(z) M(z)^* = \tilde{M}(z) M(z)^{-1} = 1$, in view of (4.14).

This means that F extends to a well-defined operator in a neighbourhood of γ , and we get

Proposition 4.1. *Let $P, \tilde{P}$ be two h -pseudodifferential operators with closed simple trajectories $\gamma \subset p^{-1}(0)$, $\tilde{\gamma} \subset \tilde{p}^{-1}(0)$ having the same period. Let $m_0 \in \gamma$, $\tilde{m}_0 \in \tilde{\gamma}$, and assume that the monodromy operators*

$$\begin{aligned} \mathcal{M}(z) &: \ker_{m_0}(P - z) \longrightarrow \ker_{m_0}(P - z) \\ \tilde{\mathcal{M}}(z) &: \ker_{\tilde{m}_0}(\tilde{P} - z) \longrightarrow \ker_{\tilde{m}_0}(\tilde{P} - z) \end{aligned}$$

are conjugate as in (4.12), via an h -Fourier integral operator $\mathcal{U}(z)$ depending smoothly on z . Then there exists a unitary Fourier integral operator F , microlocally defined in a neighbourhood of γ with an associated canonical transformation $\kappa_F : \text{neigh}(\gamma) \rightarrow \text{neigh}(\tilde{\gamma})$, such that $\tilde{P}F = FP$.

□

Using Theorem 3 we will recover the infinitesimal quantum Birkhoff normal form of $\mathcal{M}(z)$ at $z = 0$. From the proof of Proposition 4.1 it is clear that the conclusions remain valid to infinite order at γ and $z = 0$, if that assumption is made on the conjugation. Hence we obtain

Proposition 4.2. *Suppose that in Proposition 4.1 we only assume that*

$$\tilde{\mathcal{M}}(z) \equiv \mathcal{U}(z) \mathcal{M}(z) \mathcal{U}(z)^{-1}$$

to infinite order at $m = m_0$, $z = 0$ in the sense of (2.12). Then there exists a unitary Fourier integral operator F , microlocally defined in a neighbourhood of γ with an associated canonical transformation $\kappa_F : \text{neigh}(\gamma) \rightarrow \text{neigh}(\tilde{\gamma})$, such that

$$\tilde{P}F = FP, \quad \text{to infinite order along } \gamma.$$

□

5. APPLICATIONS TO INVERSE PROBLEMS

We start with a simple lemma motivated by (2.3) and (2.6):

Lemma 5.1. *Let $u(z, h)$ be a semiclassical family of functions defined near $z = 0$,*

$$u(z, h) = e^{iI(z)/h} \sum_{j=0}^{\infty} a_j(z) h^j, \quad I, a_j \in C^\infty, \quad a_0(0) \neq 0, \quad \text{Im } I(z) = 0.$$

Suppose that $\mathcal{V} \subset C_c^\infty(\mathbb{R})$ is a subspace from which we can pick an element having any prescribed finite Taylor polynomial at $I'(0)$. If for any $g \in \mathcal{V}$ we know $b_j(g)$ where

$$J(g, u) = h^{-1} \int \hat{g}(z/h) \chi(z) u(z, h) dz = e^{iI(0)/h} \sum_{k=0}^{\infty} b_k(g) h^k, \quad \chi \in C_c^\infty(\mathbb{R}), \quad \chi \equiv 1 \text{ near } 0,$$

then the distributions b_k determine the $I^{(l)}(0)$, and $a_j^{(l)}(0)$, for all l and j .

Proof. We first observe that if $a_j(z) = \mathcal{O}(z^\infty)$ then they contribute $\mathcal{O}(h^\infty)$. Hence in considering the expansion of $J(g, u)$ we can replace I and a_j 's by their formal power expansions:

$$I(z) = I_0 + I_1 z + \sum_{k=1}^{\infty} I_{k+1} z^{k+1}, \quad a_j(z) = \sum_{k=0}^{\infty} a_{jk} z^k,$$

and our task is the recovery of I_k 's and a_{jk} 's. Making a change of variables $\zeta = z/h$ we write $J(g, u)$ as the formal integral

$$e^{I_0/h} \int_{\mathbb{R}} \hat{g}(\zeta) e^{iI_1 \zeta} e^{i \sum_{j=1}^{\infty} h^j I_{j+1} \zeta^{j+1}} \left(\sum_{j,k \geq 0} a_{jk} \zeta^k h^{k+j} \right) d\zeta,$$

where we can neglect $\chi(h\zeta)$, as $\chi(z) \equiv 1$ near $z = 0$. Just as in the proof of Theorem 3 we now see that the coefficient of h^{p+1} in the expansion is given by

$$e^{I_0/h} h \int \hat{g}(\zeta) e^{iI_1 \zeta} \left(\sum_{l+k=p} a_{lk} \zeta^l + i a_{00} I_{p+1} \zeta^{p+1} + H_m(a, I) \right),$$

where $H_m(a, I)$ depends only on a_{lk} with $l + k < p$, and I_l with $l < p + 1$. Since we assumed that $a_{00} = a_0(0) \neq 0$, and that we can find $g \in \mathcal{V}$ with any prescribed finite Taylor polynomial at I_1 , the coefficients of I and a can be recovered from $b_j(g)$'s, $g \in \mathcal{V}$. $\square$

Using Lemma 5.1, and [17, Theorem 2, Proposition 7.5] (see (2.3) (2.6) above) we obtain the following consequence of Theorem 3:

Theorem 4. *Suppose that $P(z)$ satisfies the assumptions needed for (2.3) and is classical in the sense of (2.5). Suppose that in addition the eigenvalues of the linear Poincaré map of $\gamma(0)$ satisfy (2.9). Then the coefficients in the asymptotic expansion of*

$$\frac{1}{\pi} \text{tr} \int f(z/h) \bar{\partial}_z [\tilde{\chi}(z) \partial_z P(z) P(z)^{-1}] A\mathcal{L}(dz) = \sum_{k \in \mathbb{Z} \setminus \{0\}} e^{ikI/h} \sum_{j=0}^{\infty} a_{j,k}(f) h^j,$$

determine the infinitesimal quantum Birkhoff normal form of the quantum monodromy operator $M(z)$ at $z = 0$. In particular the Birkhoff normal form of the Poincaré map of $\gamma(0)$ is determined. Here the notation is the same as in (2.3), and by the coefficients we mean the distributions, $f \mapsto a_j(f)$, $\hat{f} \in C_c^\infty(\mathbb{R} \setminus \{0\})$.

Proof. We first see that the non-degeneracy of $\gamma(z)$ for z close to 0 shows that (2.6) holds. We apply (2.3) and integrate by parts:

$$\begin{aligned} & -\frac{1}{2\pi i} \sum_{-1 \neq k = -N-1}^{N-1} \operatorname{tr} \int_{\mathbb{R}} f(z/h) M(z, h)^k \frac{d}{dz} M(z, h) \chi(z) dz = \\ & h^{-1} \frac{1}{2\pi i} \sum_{-1 \neq k = -N-1}^{N-1} \operatorname{tr} \int_{\mathbb{R}} f'(z/h) (k+1)^{-1} M(z, h)^{k+1} \chi(z) + \mathcal{O}(h^\infty), \end{aligned}$$

where $\chi \equiv 1$ near 0 and hence the contribution of χ' is negligible: $f(z/h) = \mathcal{O}(h^\infty)$ for $|z| > \epsilon > 0$. The assumptions of Lemma 5.1 with $\hat{g} = f'$ are satisfied and hence we obtain expansions of $\operatorname{tr} M(z, h)^k$, for any k . Now we can apply Theorem 3 with $U(z) = M(z, h)$ and the proof is completed. $\square$

When we combine this result with Proposition 4.2 we obtain infinitesimal information about $P(z)$ at γ :

Corollary 5.2. *Suppose that the assumptions of Theorem 4 are satisfied and $P(z) = A(z)^*(P - z)A(z)$, where $A(z)$ is elliptic near γ . Then, up to conjugation by elliptic h -Fourier integral operators, in the sense of Proposition 4.2, P is determined to infinite order at γ .*

When specialized to the classical, or high energy, case, we obtain

Corollary 5.3. *Let M be a compact $\mathcal{C}^\infty$ manifold and let P be a positive, self-adjoint elliptic classical pseudo-differential operator acting on $\mathcal{C}^\infty(M, \Omega_M^{\frac{1}{2}})$. Suppose that $\gamma \subset p^{-1}(1)$ is a simple non-degenerate orbit of the classical flow of the principal symbol of P , and P_γ is its linear Poincaré map. If (2.9) holds for the eigenvalues of P_γ (2.8), and no other orbit in $p^{-1}(1)$ has the period which is a multiple of the period of γ , then the spectrum of P determines the Birkhoff normal form of the Poincaré first return map.*

Proof. Suppose that the order of P is m . We define a semi-classical operator, $P(h)$, by putting, $P(h) = h^m P(x, D_x) - 1$, so that the (semi-classical) principal symbol of $P(h)$ is the same as the (classical) principal symbol of P . The non-degeneracy assumption for $\gamma \subset \{p = 1\}$ implies that the assumptions of Theorem 4 are satisfied, with $P(z) = P(h) - z$. The simplicity of γ then shows that for f 's with $\operatorname{supp} \hat{f}$ compact, and sufficiently close to $\mathbb{Z}T_\gamma$, where T_γ is the (primitive) period of γ on $\{p = 1\}$, we have

$$(5.1) \quad \operatorname{tr} f(P(h)/h) \chi(P(h)) A = \operatorname{tr} f(P(h)/h) \chi(P(h)) + \mathcal{O}(h^\infty).$$

The left hand side is a (semi-classical) spectral invariant and hence the result follows from Theorem 4. $\square$

We leave it to the reader to formulate the corresponding corollary in the more general semi-classical case.

In the case of manifolds with boundaries we consider, for the sake of simplicity, the case of second order differential operators only:

Theorem 5. *Suppose that (M, g) is a Riemannian manifold with boundary, and that Δ_g is its Dirichlet or Neumann Laplacian.*

Let γ be a simple non-degenerate orbit of the classical billiard flow of g . If all the intersections of γ with ∂M are transversal, (2.9) holds for eigenvalues of the Poincaré map of γ (2.8), and the lengths of other closed orbits of the generalized billiard flow do not accumulate at a multiple of the period of γ , then the spectrum of Δ_g determines the Birkhoff normal form of the Poincaré first return map of γ .

Outline of the proof: First we need to indicate why the results of [17] can be applied in this situation. For that we will check that local solutions can be microlocally defined near a trajectory γ . Let p be the symbol of $D_t^2 - \Delta_g$ and q the defining function of ∂M . The transversality condition on γ means that $H_q p \neq 0$ at the boundary. We first recall (see for instance [10, Sect.24.2]) that we can find coordinates in which, near a point in ∂M , $q = x_1$, and $p(x, \xi) = \xi_1^2 - r(x, \xi')$, $\xi' = (\xi_2, \dots, \xi_{m+1})$, $m = \dim M$. Solving the boundary problem microlocally is equivalent to a hyperbolic problem and hence we have a solution. For localization needed in (2.3) we can use, near the boundary, an operator $A = A(x, hD_{x'})$.²

A more subtle argument is needed for showing that (5.1) holds, as that involves results on propagation of singularities for boundary value problems – see [10, Sect.24.5] and references given there. These results provide the foundation for the trace formula of Guillemin and Melrose [8] (see also [15]), which among other things shows that $\text{singsupp}(\text{tr} \cos(t\sqrt{-\Delta_g})) \subset \overline{\mathcal{L}}$, where $\mathcal{L}$ is the set of length of closed orbits of the billiard flow. It also follows from [8] and [15] that if $A(x, \langle D_t \rangle^{-1} D_x)$ is an operator which provides a localization to a neighbourhood of the bicharacteristic strip of γ , then

$$(5.2) \quad \text{tr} \cos(t\sqrt{-\Delta_g}) - \text{tr} \cos(t\sqrt{-\Delta_g}) A(x, \langle D_t \rangle^{-1} D_x) \in \mathcal{C}^\infty(\mathbb{R}), \quad \text{for } t \text{ near } kT_\gamma,$$

that is, the singularities of the trace of $\cos(t\sqrt{-\Delta_g})$ at kT_γ come from a small neighbourhood of γ (depending on k). As stated above we only need tangential pseudo-differential operators to localize near γ .

A straightforward translation to the semi-classical setting (see for instance [23]) gives (5.1). In fact, by standard oscillatory testing, $A(x, \langle D_t \rangle^{-1} D_x) e^{it/h} = e^{it/h} \tilde{A}(x, hD_x; h)$, where $\tilde{A}$ has the semi-classical principal symbol given by $A(x, \xi)$. This shows that for $\hat{f} \in \mathcal{C}_c^\infty(\mathbb{R})$ even,

$$\text{tr} f((h\sqrt{-\Delta_g} - 1)/h) \tilde{A}(x, hD_x, h) = \frac{1}{2\pi} \int \text{tr} \left(A(x, \langle D_t \rangle^{-1} D_x) \cos t\sqrt{-\Delta_g} \right) \hat{f}(t) e^{it/h} dt,$$

and thus (5.2) follows from (5.1). $\square$

By combining Theorem 4, or rather Corollary 5.3, with Proposition 4.2 or with the geometric arguments of [7, §2] we obtain the *Weinstein conjecture* first proved by Guillemin [7] in the case of elliptic trajectories, and by Zelditch [19],[20] in general. It involves the concept of the *Birkhoff normal form for the Hamiltonian* – [7, Theorem 1.1]. As shown by Françoise and Guillemin [4] and Guillemin [7, §2], and as can be also deduced from Sect.4 above, that normal form is essentially equivalent to the normal form of the Poincaré map.

Corollary 5.4. *Let (M, g) be a compact Riemannian manifold without boundary, and γ a simple non-degenerate geodesic on M , with the spectrum of its linear Poincaré map satisfying (2.9). Then the spectrum of the Laplacian, Δ_g , determines the Birkhoff normal form of γ .*

In the special case of symmetric domains in the plane, we can apply Theorem 5 and an observation of Colin de Verdière [2, §4] to recover a result of Zelditch [21]³:

Corollary 5.5. *Suppose that $\Omega \subset \mathbb{R}^2$ is a bounded domain, $\partial\Omega$ is real analytic, and that Ω is symmetric with respect to the x and y axes. Suppose also that Ω has a simple non-degenerate bouncing ball orbit along one of the axes. Then the spectrum of the Dirichlet (or Neumann) Laplacian determines Ω among all such domains.*

²Alternatively, we could have reduced the problem to the boundary, and considered the monodromy operator there.

³Melrose [14] has announced the same result, sketching a yet different proof. Recently, by considering the full quantum Birkhoff normal form, Zelditch [22] succeeded in removing one of the symmetry conditions

REFERENCES

- [1] N.W. Ashcroft and N.D. Mermin, *Solid State Physics* Saunders College Publishing, Philadelphia, 1976.
- [2] Y. Colin de Verdière, *Sur les longueurs de trajectoires périodiques d'un billard*, in P. Dazord and N. Desolmes-Moulis (eds.) *Géométrie Symplectique et de Contact, South Rhone seminar on geometry, III (Lyon, 1983)*, Herman, Paris, 1984, 122-139.
- [3] M. Dimassi and J. Sjöstrand, *Spectral Asymptotics in the semi-classical limit*, Cambridge University Press, 1999.
- [4] J.P. Francoise and V. Guillemin, *On the period spectrum of a symplectic mapping*, J. Funct. Anal. **115**(1993), 391-418.
- [5] D. Fried, *Cyclic resultants of reciprocal polynomials*, in *Holomorphic Dynamics (Mexico 1986)*, Lecture Notes in Math. **1345**, Springer Verlag, 1988, 124-128.
- [6] V. Guillemin, *Wave trace invariants and a theorem of Zelditch*, Int. Math. Res. Not. **12**(1993), 303-308.
- [7] V. Guillemin, *Wave trace invariants*, Duke Math. J. **83**(1996), 287-352.
- [8] V. Guillemin and R.B. Melrose, *The Poisson formula for manifolds with boundary*, Adv. in Math. **32**(1979), 204-232.
- [9] B. Helffer and J. Sjöstrand, *On the diaphanization and the de Haas-van Alphen effect* Ann. Inst. H. Poincaré Phys. Théor. **52**(1990), no. 4, 303-375.
- [10] L. Hörmander, *The Analysis of Linear Partial Differential Operators, vol. III, IV*, Springer Verlag, 1985.
- [11] A. Iantchenko, *La forme normale de Birkhoff pour un opérateur intégral de Fourier*, Asympt. Anal. **17**(1998), 71-92.
- [12] A. Iantchenko and J. Sjöstrand, *Birkhoff normal forms for Fourier integral operators II*, preprint, 2001. <http://xxx.lanl.gov/abs/math.SP/0111134>
- [13] M.P. Jacobson, C. Jung, H.S. Taylor, and R.W. Field, *State-by-state assignment of the bending spectrum of acetylene at 15 000 cm⁻¹: A case study of quantum-classical correspondence*, J. Chem. Phys. **111**(2)(1999), 600-618.
- [14] R.B. Melrose, *Inverse spectral problem for planar domains (joint work with V.W. Guillemin)*, Talk at Berkeley, September, 1998.
- [15] V. Petkov and L. Stoyanov, *Geometry of Reflected Rays and Inverse Spectral Problems*, John Wiley & Sons, New York 1992.
- [16] J. Sjöstrand, *Semi-excited states in non-degenerate potential wells*, Asympt. Anal. **6**(1992), 29-43.
- [17] J. Sjöstrand and M. Zworski, *Quantum monodromy and semi-classical trace formulae*, to appear in J. Math. Pure Appl. <http://xxx.lanl.gov/abs/math.SP/0108052>
- [18] R.B. Shirts and W.P. Reinhardt, *Approximate constants of motion for classically vibrational dynamics: Vague tori, semiclassical quantization, and classical intramolecular energy flow*. J. Chem. Phys. **77**(10)(1982), 5204-5216.
- [19] S. Zelditch, *Wave invariants at elliptic closed geodesics*, Geom. Funct. Anal. **7**(1997), 145-213.
- [20] S. Zelditch, *Wave invariants for non-degenerate closed geodesics*, Geom. Funct. Anal. **8**(1998), 179-217.
- [21] S. Zelditch, *Spectral determination of analytic bi-axisymmetric domains*, Geom. Funct. Anal. **10**(2000), 628-677.
- [22] S. Zelditch, *The inverse spectral problem for analytic plane domains. I: Balian-Bloch trace formula, II: domains with symmetry*, preprints 2001, <http://arXiv.org/abs/math.SP/0111077>, <http://arXiv.org/abs/math.SP/0111078>
- [23] M. Zworski, *A remark on a paper by E.B. Davies*, Proc. A.M.S. **129**(2001), 2955-2957.

MALMÖ HÖGSKOLA, TEKNIK OCH SAMHÄLLE, SE-20506, MALMÖ, SWEDEN

E-mail address: ai@ts.mah.se

CENTRE DE MATHÉMATIQUES, ÉCOLE POLYTECHNIQUE, FR-91128, PALAISEAU CEDEX, FRANCE, AND UMR 7640 OF CNRS

E-mail address: johannes@math.polytechnique.fr

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CA 94720, USA

E-mail address: zworski@math.berkeley.edu